

## No penicillin allergy

**First-line: 7 days, PPI twice daily**  
**plus** amoxicillin 1 g b.d.  
**plus** either clarithromycin 500 mg b.d. **or**  
metronidazole 400 mg b.d.

**Ongoing symptoms** after first-line

**Second-line: 7 days, PPI twice daily**  
**plus** amoxicillin 1 g b.d.  
**plus** second antibiotic not used  
in first-line, either clarithromycin  
500 mg b.d. **or** metronidazole  
400 mg b.d.

**Ongoing symptoms** after first-line **and**  
previous exposure to metronidazole **and**  
clarithromycin

**Second-line: 7 days, PPI twice daily**  
**plus** amoxicillin 1 g b.d.  
**plus** second antibiotic, either tetracycline  
hydrochloride 500 mg q.d.s **or**  
levofloxacin 250 mg b.d.

## Penicillin allergy

**First-line: 7 days, PPI twice daily**  
**plus** clarithromycin 500 mg b.d.  
**plus** metronidazole 400 mg b.d.

First-line with previous clarithromycin  
exposure **or**  
Second-line with previous levofloxacin  
exposure

**7 days, PPI twice daily**  
**plus** bismuth subsalicylate 525 mg q.d.s  
**or** tripotassium dicitratobismuthate  
240 mg q.d.s  
**plus** tetracycline hydrochloride  
500 mg q.d.s  
**plus** metronidazole 400 mg b.d.

**Ongoing symptoms** after first-line **and**  
**no** previous exposure to levofloxacin

**Second-line: 7 days, PPI twice daily**  
**plus** metronidazole 400 mg b.d.  
**plus** levofloxacin 250 mg b.d.